7. REFERENCES

1.AbdulrahmanS,Md.Afroz Ahmad (2015), Hepatoprotective and antioxidant activity of Tragia involucrata root extracts against CCl4 induced hepatotoxicity in rats. Scholars research libraray, vol7 (5),pp 146-152.

2.AdigaG ,Grover JK (2001), "Extracts of Benincasa hipsida prevents development of experimental ulcers". Journal of Ethnopharmacology, vol7(8), pp159-64.

3.Ajitkar, Bijoykumarchoudhary, swarupkumar (2006) , a process for the preparation of 5-methyl-isoxazole-3-methyl carboxylate (a novel hypoglycaemic compound with hypolipidaemic and antioxidant activities) from the plant tragia involucrata, The patent office journal p15335.

4. A. K. Dhara, NagChaudhuri,S. Pal (2002), Psychopharmacological studies on Tragia involucrata root extract. Phytotherapy Research, vol16 (4), pp 326–330.

5. A.KDhara, Nag Chaudhuri,Pal S , SubaV, Sen T (2000). Preliminary studies on the anti-inflammatory and analgesic activity of the methanolic fraction of the root extract of TragiaInvolucrataLinn.Journal of ethnopharmacology, vol72 (12),pp 265-268.

6. Benson K.Mathai,GanapathiG.varma,,Kunthal Das(2014),Evaluation of antiepileptic activity of methanolic leaves extract of Tragia involucrate Linn. In mice. International letters of natural sciences, vol17 (2), pp167-17.

7. Debashisha Panda (2012) Phytochemical Examination and Antimicrobial Activity of Solvent Extracts and Isolatation of Compounds from Roots of Tragia involucrata.International Journal of Pharmaceutical Sciences and Drug Research , vol 4(1), pp44-48.

8. Deyet (2011), "Alstonia scholaris R.Br. (Apocynaceae): Phytochemistry and pharmacology: A concise review". Journal of Applied Pharmaceutical Sciences, vol 1(6), pp51-57.

9. G. Don Byernawita (2008) Bioassay-guided fractionation and identification of antioxidant and antimicrobial compounds from callistemon viminalis.Journal of pharmacology and toxicology, vol 3(2), pp10-14.

10. G.Chandrashekhar Joshi and M.Gopal (2011), Antifertility activity of hexane and ethyl acetate extracts of aerial parts of tragia involucrata Linn. Journal of pharmacology and toxicology, vol4 (6), pp. 11-19

11. H.G,Vogel,W.HVogel (1997), Drug discovery Evaluation, New york, Springer velag Berlin Heidelberg, pp. 418

12. Jain MS ,Rachchh MA (2008), "Gastro-protective effect of Benincas ahipsida fruit extract". Indian Journal of Pharmacology, vol40(6), pp271-75.

13. Janakiraman N and Johnson M,SahayaSathish S (2012), Phytochemical Analysis of Vitexaltissima L. using UV-VIS, FTIR and GC-MS. International Journal of Pharmaceutical Sciences and Drug Research , vol4(1), pp 56-62

14. IgnacimuthuS, Samy RP (2000), Anti inflammatory effect of different extracts of tragia involucrata for carageenan induced hind paw oedema in wistar albino rats.journal of medical and aromatic plantsciences, vol22(1), pp 34-35.

15. IndiraBalachandran, Sivaraman (1994), Ayurvedi Drugs and Their Plant Source.

Oxford and IBH publishing Co.PvtLtd, pp.147.

16. K.Benoy, K.Hemamalini1, S.M. ShantaKumar, S. Satyanarayana N.VenkatRao (2007), Pharmacological evaluation of root extracts of tragia involucrata. Pharmacologyonline, vol 2(2), pp 236-244

17. K.Benoy, K.Hemamalini, S.M. Satyanarayan, N.VenkatRao(2007), Anti-diabetic activity of root extracts of Tragia involucrata. Pharmacologyonline, vol2 (4) pp 203-217.

18. Kokate CK.(1994), Practical Pharmacognosy, VallabhPrakashan ,4th edn, pp107-111.

19. Kaushik M, Kiran D MurtyK,Paliwal D (2011), “Preliminary and pharmacological evaluation profile of Ficus religiosa: An overview”. Pharmacologyonline, vol3(1), pp387-395.

20. Khandelwal KR. Practical Pharmacognosy (2000), Techniques and experiments. 2nd edn.Nirali Prakashan, pp 146-170

21. M.Alimuzzaman, MuniruddinAhmed (2005), Analgesic Activity of Tragia involucrata. Dhaka University journal of pharmaceutical sciences, Vol.4 (1) pp.10

22. M. Khazaei, H. Salehi (2006), Protective effect of Falcaria vulgaris extract on ethanol induced gastric ulcer in rat. Iranian Journal of Pharmacology and Therapeutics, vol 5 (2), pp1–4.

23. M Madhuri,Mohammad Saleem TS, Pradeep KR, Priyanka N, Sravanti U, Sarala K(2011), Anti-ulcerogenic activity of aqueous extract of Annona squamosa Linn. International Journal of Research in Phytochemistry and Pharmacology,vol 2(3), pp 157-159.

24. Mahobia N, Patel R, Talaviya H Upwar N, Waseem N (2010), “Analgesic and antipyretic activities of Momordica charantia Linn. fruits”. Journal of Advance Pahrmaceutical and Technology and Research, vol1 (4), pp414-18.

25. Mohd. AzeemuddinMukhram et al. (2011) Gastrokinetic and Gastroprotective Activity of Gasex (an Ayurvedic preparation of The Himalaya Drug Company).journal of ayurvedicmedicines, vol 4(7), pp. 23-26.

25. M. LingaRao, N. Savithramma and D. Suhrulatha (2011), Screening of Medicinal Plants for Secondary Metabolites, Middle-East Journal of Scientific Research vol 8 (3), pp579-584.

26. María Elena Sánchez-Mendoza et al (2011), Bioassay-Guided Isolation of an Anti-Ulcer Compound,Tagitinin C, from Tithoniadiversifolia. Journal molecules, vol16, pp665-674.

27. Maria Elena San (2015),Gastroprotective Activity of Methyleugenol from Peperomia hispidula on Ethanol-Induced Gastric Lesions in Rats. International Journal of Pharmacology, vol11 (7), pp 697-704.

28. Muthuraman meenakshisundaram et al (2009), Chemotaxonomic studies on Tragiainvolucrata Linn, International journal of Biol. Chem. Sci. 3(5), pp. 927-933.

29. M.Rama Devi et al (2010), Anti gasric ulcer activity of plectranthus amboinicus(lour) in wistar albino rats. Journal of chemical and pharmaceutical research, 2(3), pp. 374-380

30. N.VenkatRao et al (2007),Anti-diabetic activity of root extracts of Tragia involucrata. Pharmacology online vol 2, pp 203-217.

31. N.VenkatRao et al (2007) pharmacological evaluation of root extracts of Tragia involucrata, Pharmacologyonline, vol 2(2), pp236-244 .

32. OECD: Guideline 420-Annex 3,Test guidelines for Toxicity studies.

33.Palani R ,SahayaSathish S, Thamizharasi T ,Vijayakanth P, and Vimala.A (2013), antimicrobial and phytochemical screening of Tragia involucrata Usinguv-vis and FT-IR.International journal of research in engineering and bioscience,vol 2(I)), pp82-90.

34. Parmar NS, Parmar S (1998), Anti-ulcer potential of flavonoids. Indian Journal of Physiology and Pharmacology, vol 4(2), pp343-35.

35. RamanKutty C ,Warrier PK, Nambiar VP(1992) Indian Medicinal Plants: A compendium of 500 species, Orient Longman Publishers.vol5(7), pp 23.

36. Samy R.P (2006), Purification of antibacterial agents from Tragia involucrata--a popular tribal medicine for wound healing. Journal of Ethnopharmacology, vol107(1), pp 99-106.

36. SaheedSabiu (2015), Indomethacin-induced gastric ulceration in rats, Protective roles of Spondias mombin and Ficus exasperate. Toxicology report, vol 2, pp 261-267.

37. Syeda JeelaniBasri (2014) K.N.Jayaveera,G.V.Subba Reddy**,** Astudy on phytochemical and chromatographic assay on Tragia involucrata, world journal of pharmacy and pharmaceutical sciences , vol3(7), pp1667-1670.

38. S. Mohamed farook et al (2011) Antidiabetic and hypolipidemic potential of tragia involucrate in streptozotocin-nicotinamide induced type ii diabetic rats. International Journal of Pharmacy and Pharmaceutical Sciences,vol 3(4), pp 20-21.

39. Thillaivanan.s et al (2015) A Review on "Kapa SuraKudineer"-A Siddha Formulary Prediction for Swine Flu, International Journal of Pharmaceutical Sciences and Drug Research ,vol7(5), pp376-383.

40. Vithyarathinam PF (1996), Indian medicinal plant: A compendium of 500 species. Orientlongman ,vol.5, pp 304.

41. ZanariahBintiUjang (2013), Bioassay guided Fractionation and Purification of Natural Bioactive molecules from Alpinia conchigera. Journal of pharmacology and toxicology, vol 4(7), pp12-17.

.